Downloaded from www.mycourses.co.za

higher education & training

Department:
Higher Education and Training
REPUBLIC OF SOUTH AFRICA

NATIONAL CERTIFICATE

MATHEMATICS N5

(16030175)

2 April 2020 (X-paper)
09:00–12:00

Scientific calculators may be used.

This question paper consists of 5 pages and a formula sheet of 4 pages.

068Q1A2002

Downloaded from www.mycourses.co.za

Downloaded from www.mycourses.co.za

DEPARTMENT OF HIGHER EDUCATION AND TRAINING
REPUBLIC OF SOUTH AFRICA

NATIONAL CERTIFICATE
MATHEMATICS N5
TIME: 3 HOURS
MARKS: 100

INSTRUCTIONS AND INFORMATION

1. Answer all the questions.
2. Read all the questions carefully.
3. Number the answers according to the numbering system used in this question paper.
4. Show all intermediate steps and simplify where possible.
5. All final answers must be rounded off to THREE decimals.
6. Questions may be answered in any order, but subsections of questions must be kept together.
7. Sketches must be large, neat and fully labelled
8. Write neatly and legibly.

Downloaded from www.mycourses.co.za

Downloaded from www.mycourses.co.za

QUESTION 1

1.1 Determine the following limits:

1.1.1 $\lim_{x \to \frac{1}{9}} \frac{9x - 1}{3\sqrt{x} - 1}$ (3)

1.1.2 $\lim_{x \to 0} (\operatorname{cosec} x - \cot x)$ (4)

1.2 Determine the value(s) of x for which $f(x)$ is discontinuous:

$$f(x) = \frac{x^2 + 4x - 12}{x^2 - 2x}$$

(2)
[9]

QUESTION 2

2.1 Derive a formula to determine $\frac{dy}{dx}$; if $y = arc \cot x$. (2)

2.2 Determine $\frac{dy}{dx}$ in each of the following cases (simplification NOT required):

2.2.1 $y = ln[\cos^5(3x^4)]$ (4)

2.2.2 $y = \left(\frac{8x - x^6}{x^3}\right)^{-\frac{4}{5}}$ (4)

2.2.3 $y = \sqrt{1 + \sqrt{1 + x^2}}$ (3)

2.3 Calculate $\frac{dy}{dx}$ if $y = \left(1 + \frac{1}{x}\right)^x$ with aid of logarithmic differentiation. (4)

2.4 Given the implicit function $25x^2 + 9y^2 - 70x - 30y + 49 = 0$

2.4.1 Determine $\frac{dy}{dx}$ (3)

2.4.2 Determine the equation of the tangent to the graph at the point (2;3). (3)
[23]

Downloaded from www.mycourses.co.za

Downloaded from www.mycourses.co.za

QUESTION 3

3.1 Given: $f(x) = x^3 - 3x + 1$

3.1.1 Determine the coordinates of the turning points of $f(x)$ (2)

3.1.2 Draw up a table of x and $f(x)$, where x is ranging from $x = -2$ to $x = 2$. (3)

3.1.3 Draw a neat graph of $f(x)$ between these values showing the turning points on it. (2)

3.1.4 Use the table and the graph to estimate a value for the best root between $x = 1$ and $x = 2$ of the equation $x^3 - 3x + 1 = 0$ and then use Taylor's/Newton's method TWICE to determine a better approximation of this root. (Root correct to THREE decimal figures) (4)

3.2 Water is being poured into a conical reservoir at a rate of $\pi m^3/s$. The reservoir has a radius of 6 metres across the top and a height of 12 metres.

At what rate is the depth of water increasing when the depth is 6 metres?

HINT: $V = \frac{1}{3}\pi r^2 h$

(4)
[15]

QUESTION 4

4.1 Determine $\int y\,dx$ in each of the following cases:

4.1.1 $y = \dfrac{4}{\sec x e^{\sin x}}$ (3)

4.1.2 $y = \dfrac{(\cos^{-1} 2x)^4}{\sqrt{1 - 4x^2}}$ (3)

4.1.3 $y = \dfrac{x^3 + 2x^2 + 9x - 17}{x + 4}$ (5)

4.1.4 $y = \cos 3\pi x . \cos \pi x$ (3)

4.1.5 $y = x\sqrt{1 + 2x}$ (4)

Downloaded from www.mycourses.co.za

Downloaded from www.mycourses.co.za

4.2 Determine $\int y\,dx$ by resolving the integral into partial fractions:

$$y = \frac{3x}{(2x+1)(x+4)}$$

(5)
[23]

QUESTION 5

5.1 Determine $\int_0^\infty e^{-st}.f(t)dt$ if $f(t) = \frac{t}{\pi}$ (5)

5.2 Given : $y = x + 1,\ x = 0$ and $y = 2x^2$

5.2.1 Calculate the coordinates of the points of intersection. (2)

5.2.2 Make a neat sketch to show the enclosed area, the representative strip and the point of intersection. (2)

5.2.3 Calculate the magnitude of the area in QUESTION 5.2.2. (3)

5.2.4 Calculate the volume of the solid of revolution formed when the area in QUESTION 5.2.2 is rotated about the x-axis. (4)

5.3 Calculate the second moment of area of a rectangular lamina with sides 6 cm × 3 cm and about a 3 cm side. (4)
[20]

QUESTION 6

6.1 Determine the particular solution of $\frac{dy}{dx} = \frac{x(e^{x^2} + 2)}{6y^2}, y(0) = 1$ (4)

6.2 Determine the particular solution of $x^2.\frac{d^2y}{dx^2} = x^6 + x^3 - 2x^2$ if $\frac{dy}{dx} = \frac{3}{5}, x = 1$ and $y = \frac{1}{5}$

(6)
[10]

TOTAL: 100

Downloaded from www.mycourses.co.za

Copyright reserved

Downloaded from www.mycourses.co.za

MATHEMATICS N5

FORMULA SHEET

Any applicable formula may also be used

TRIGONOMETRY

$\sin^2 x + \cos^2 x = 1$
$1 + \tan^2 x = \sec^2 x$
$1 + \cot^2 x = \text{cosec}^2 x$
$\sin 2A = 2 \sin A \cos A$
$\cos 2A = \cos^2 A - \sin^2 A$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$$
$\sin^2 A = {}^1/_2 - {}^1/_2 \cos 2A$
$\cos^2 A = {}^1/_2 + {}^1/_2 \cos 2A$

$\sin(A \pm B) = \sin A \cos B \pm \sin B \cos A$
$\cos(A \pm B) = \cos A \cos B \pm \sin A \sin B$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \pm \tan A \tan B}$$

$\sin A \cos B = {}^1/_2 [\sin(A + B) + \sin(A - B)]$
$\cos A \sin B = {}^1/_2 [\sin(A + B) - \sin(A - B)]$
$\cos A \cos B = {}^1/_2 [\cos(A + B) + \cos(A - B)]$
$\sin A \sin B = {}^1/_2 [\cos(A - B) - \cos(A + B)]$

$$\tan x = \frac{\sin x}{\cos x}; \sin x = \frac{1}{cosec\ x}; \cos x = \frac{1}{\sec x}$$

BINOMIAL THEOREM

$$(x + h)^n = x^n + \mathrm{n}x^{n-1}\, h + \frac{n(n-1)}{2!} x^{n-2}\, h^2 + \cdots$$

DIFFERENTIATION

$$e = -\frac{f(a)}{f'(a)}$$
$r = a + e$

PRODUCT RULE

$\mathrm{y} = \mathrm{u}(x) \,.\, v(x)$
$$\frac{dy}{dx} = u \,.\, \frac{dv}{dx} + v \,.\, \frac{du}{dx}$$
$= u \,.\, v' + v \,.\, u'$

Downloaded from www.mycourses.co.za

Copyright reserved

Downloaded from www.mycourses.co.za

QUOTIENT RULE

$$y = \frac{u(x)}{v(x)}$$

$$\frac{dy}{dx} = \frac{v\,.\,\frac{du}{dx} - u\,.\,\frac{dv}{dx}}{v^2}$$

$$\frac{v\,.\,u' - u\,.\,v'}{v^2}$$

CHAIN RULE

$$y = f\big(u(x)\big)$$

$$\frac{dy}{dx} = \frac{dy}{du}\,.\,\frac{du}{dx}$$

$f(x)$	$\frac{d}{dx}f(x)$	$\int f(x)dx$
ax^n	nax^{n-1}	$\frac{ax^{n+1}}{n+1} + c$
a	0	$ax + c$
e^x	e^x	$e^x + c$
a^x	$a^x.lna$	$\frac{a^x}{lna} + c$
$\log_e x$	$\frac{1}{x}$	___
$\log_a x$	$\frac{1}{xlna}$	___
$\sin x$	$\cos x$	$-\cos x + c$
$\cos x$	$-\sin x$	$\sin x + c$
$\tan x$	$\sec^2 x$	$ln(\sec x) + c$
$\cot x$	$-\operatorname{cosec}^2 x$	$ln(\sin x) + c$
$\sec x$	$\sec x\,.\tan x$	$ln[\sec x + \tan x] +$
c		
$\operatorname{cosec} x$	$-\operatorname{cosec} x\,.\cos x$	$ln(\operatorname{cosec} x +$
$\cot x) + c$		
$\sin^{-1} x$	$\frac{1}{\sqrt{1-x^2}}$	___
$\cos^{-1} x$	$\frac{-1}{\sqrt{1-x^2}}$	___
$\tan^{-1} x$	$\frac{1}{1+x^2}$	___
$\cot^{-1} x$	$\frac{-1}{1+x^2}$	___
$\sec^{-1} x$	$\frac{1}{\sqrt{x^2-1}}$	___
$\operatorname{cosec}^{-1} x$	$\frac{-1}{x\sqrt{x^2-1}}$	___
$f(x)$	$\frac{d}{dx}f(x)$	$\int f(x)dx$

Downloaded from www.mycourses.co.za

Copyright reserved

Downloaded from www.mycourses.co.za

$\frac{1}{\sqrt{a^2-x^2}}$	—	$\sin^{-1}\left(\frac{x}{a}\right) + c$
$\frac{1}{a^2+x^2}$	—	$\frac{1}{a}\tan^{-1}\left(\frac{x}{a}\right) + c$
$\frac{1}{x\sqrt{x^2+a^2}}$	—	$\frac{1}{a}\sec^{-1}\left(\frac{x}{a}\right) + c$
$\sqrt{a^2-x^2}$	—	$\frac{a^2}{2}\sin^{-1}\left(\frac{x}{a}\right) +$
$\frac{x}{2}\sqrt{a^2-x^2} + c$		
$\frac{1}{x^2-a^2}$	—	$\frac{1}{2a}ln\left(\frac{x-a}{x+a}\right) + c$
$\frac{1}{a^2-x^2}$	—	$\frac{1}{2a}ln\left(\frac{a+x}{a-x}\right) + c$

INTEGRATION

$$\int f(x)\, g'(x)\, dx = f(x)\, g(x) - \int f'(x) g\,(x) dx$$

$$\int \frac{f'(x)}{f(x)} dx = ln\, f(x) + c$$

$$\int [f(x)]^n\, f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c$$

$$\frac{f(x)}{(ax+b)(cx+d)} = \frac{A}{ax+b} + \frac{B}{cx+d}$$

$$\frac{f(x)}{(x+a)^n} = \frac{A}{(x+a)} + \frac{B}{(x+a)^2} + \frac{C}{(x+a)^3} + \cdots \frac{Z}{(x+a)^n}$$

APPLICATIONS OF INTEGRATION

AREAS

$$A_x = \int_a^b y dx\ ;\ A_x = \int_a^b (y_1 - y_1) dx$$

$$A_x = \int_a^b x dy\ ; A_{Y=} \int_a^b (x_1 - x_2) dy$$

VOLUMES

$$V_x = \pi \int_a^b y^2\, dx\ ;\ V_{X=\pi} \int_a^b (y_1^2 - y_2^2) dx$$

$$V_y = \pi \int_a^b x^2\, dy\ ;\ V_{y=\pi} \int_a^b (x_1^2 - x_2^2) dy$$

Downloaded from www.mycourses.co.za

Downloaded from www.mycourses.co.za

SECOND MOMENT OF AREA

$$I_x = \int_a^b r^2\, dA\ ;\ I_{y=} \int_a^b r^2\, dA$$

MOMENT OF INERTIA

$$Mass\ = density\ \times volume$$
$$M\ = pv$$
$$DEFINITION : I\ = m\, r^2$$
$$GENERAL : I\ = \int_a^b r^2 dm\ = p \int_a^b r^2 dV$$

Downloaded from www.mycourses.co.za

Copyright reserved